

## Changes on weed flora from 1976 and thirty years later in Spanish winter cereals

A. Cirujeda<sup>1</sup>, A. Mari<sup>1</sup>, J. Aibar<sup>2</sup> and C. Zaragoza<sup>1</sup>

<sup>1</sup>Unidad de Sanidad Vegetal, CITA, Avda. Montañana 930, 50059 Zaragoza, Spain

acirujeda@aragon.es, carza@aragon.es

<sup>2</sup>EPS. Universidad de Zaragoza, Ctra. de Cuarte, Km 67, 22071 Huesca, Spain. jaibar@unizar.es

In this work we compare the weed flora recorded in 1976 with that in 2005-07 in winter cereal fields in the province of Zaragoza. In 2005-07 we used a very similar methodology to that used in 1976, based on surveying the fields in a zig-zag by three persons until not founding new species and assessing the abundance of species with a visual scale. It is concluded that grass species have increased, especially *Lolium rigidum* Gaudin and *Avena* spp. The average number of species found in each field has decreased from  $9.5 \pm 2.3$  in 1976 to  $3.4 \pm 2.3$  in 2005-07. In 1976, 22 species were found in more than 30% of the fields studied, in 2005-07 only 4 species were counted. Species that have declined more drastically are *Daucus carota* L., *Scandix pecten-veneris* L., *Veronica agrestis* L., *Papaver hybridum* L., *Roemeria hybrida* (L.) DC. and *Hypochaeris procumbens* L.. However, we have found 23 species, which were not recorded in 1976, including: *Chondrilla juncea* L., *Phragmites australis* (Cav.) Trin., *Sorghum halepense* (L.) Pers., *Xanthium strumarium* L., *Kochia scoparia* (L.) Schrader, *Hordeum murinum* L.. Some are linked to minimum tillage, others are tolerant to glyphosate pre-sowing treatments (*Equisetum arvense* L., *Malva sylvestris* L.) and others may have advanced their emergence due to global warming.